

14.07.00

PCT/

00/00406



Eur päisches
Patentamt

European
Patent Office

Office eur péen
des brevets

10-009874

NL00
00406

REC'D 10 AUG 2000	
WIPO	PCT

Bescheinigung

Certificate

Attestation

4

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

99201862.2

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

I.L.C. HATTEN-HECKMAN

DEN HAAG, DEN
THE HAGUE,
LA HAYE, LE

26/06/00

4

4

4

4



**Eur päisches
Patentamt**

**Eur pean
Patent Office**

**Office eur péen
des brevets**

**Blatt 2 der Bescheinigung
Sheet 2 of the certificate
Page 2 de l'attestation**

Anmeldung Nr.:
Application no.: 99201862.2
Demande n°:

Anmeldetag:
Date of filing: 11/06/99
Date de dépôt:

Anmelder:
Applicant(s):
Demandeur(s):
Landbouwniversiteit Wageningen
6700 HB Wageningen
NETHERLANDS

Bezeichnung der Erfindung:
Title of the invention:
Titre de l'invention:

Expression in plants of starch binding domains and/or of fusion proteins containing starch binding domains

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:
State:
Pays:

Tag:
Date:
Date:

Aktenzeichen:
File no.
Numéro de dépôt:

Internationale Patentklassifikation:
International Patent classification:
Classification internationale des brevets:

C12N15/54, C12N15/62, C12N15/82, C08B30/00, A01H5/00, A01H5/10, A23L1/0522

Am Anmeldetag benannte Vertragsstaaten:
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE
Etats contractants désignés lors du dépôt:

Bemerkungen:
Remarks:
Remarques:



11-06-1999

WAGENINGEN

EP99201862.2

DESC

1

Expression in plants of starch binding domains and/or of protein fusions containing starch binding domains.

The present invention relates to methods for gene expression in plants.

5 In particular, the invention relates to the expression in plants of proteins and/or polypeptides - including enzymes - as fusions with improved affinity for starch, such as for the starch granules that may occur in plants *in vivo*, in particular in cellular organelles such as plastids

10 More in particular, the invention relates to the expression in plants of proteins and/or polypeptides as a fusion with one or more moieties or domains that have affinity for starch, such as 'starch binding domains'.

Some further aspects of the invention reside in the protein fusions thus expressed, in gene constructs that encode such fusions, and in methods for the transformation of plants using such constructs as well as in the transformed plants thus obtained.

15 The invention can in particular be used for the expression in plants *in vivo* of enzymes as fusions that become associated with the starch granules that occur in (some of the cellular organelles of) such economically important plants as potato, sweet potato, cassava, pea, taro, sago, yam, banana, and/or cereals such as maize, rice wheat, and barley. This may either facilitate the isolation of the enzymes thus expressed from
20 the plant material, as well as their further use, and/or the enzymes thus expressed may be used to alter and/or otherwise influence the starch or starch granules leading to plants with improved properties, such as the production of modified starches. These and other applications of the invention will become clear from the description hereinbelow.

25 In starch-producing plants, starch is usually synthesized/present in the form of starch granules. A number of enzymes in the plant are known to interact *in vivo* with these granules, for instance in order to build up, modify and/or degrade the starch molecules, the starch granules and/or the structure thereof. These include enzymes such as starch synthases, branching and debranching enzymes, etc., for which in general
30 reference is made to A.M. Smith, K. Denyer and C. Martin, *Annu. Rev. Plant Physiol. Mol. Biol.*, 1997, 48:67-87 and C. Martin and A.M. Smith, *The Plant Cell*, Vol.7, 971-985, July 1995.

Such enzymes or mutants thereof may also be used to produce modified starches, either in vitro or *in vivo*. The production of modified starches by plants transformed with (genes encoding) such enzymes is for instance described in DE-A-195 34 759, WO 92/14827 (in which a branching enzyme derived from potato cDNA is used) and WO 92/11376 (which describes an alternative method in which antisense DNA is used to suppress GBSS activity in plants.)

It is also known that some micro-organisms contain proteins/enzymes that can interact with (i.e. degrade, modify or convert) starch or starch granules, and some non-limiting examples thereof are mentioned below. Again, such enzymes and/or mutants thereof have been used in the art to produce modified starches in vitro and/or *in vivo*, the latter in plants transformed with (a) gene(s) encoding said enzyme or mutant. Reference is for instance made to WO 91/19808, US-A-5,349,123 and US-A-5,750,875, which describe plants transformed with bacterial amylases - and in particular cyclodextringlycosyltransferases (CGTases) - for the *in vivo* production of cyclodextrins.

Proteins/enzymes that can interact with starch, and in particular the enzymes from micro-organisms, generally contain - besides one or more catalytic domains - one or more regions that can bind to the starch and/or the starch granules. The latter are referred to in the art as "starch binding regions" or "starch binding domains". For a description of some starch binding domains that have been investigated in the art, reference is for instance made to Penninga et al., J. Biol. Chem, Vol.271, No 51, 32777-32784, 1996, and Lawson et al., J. Mol. Biol. (1994) 236, 590-600, who describe the raw starch binding domain (E-domain) of CGTase from *B.circulans*; Sorimachi et al., J. Mol. Biol. (1996), 259, 970-987 and Structure 1997, Vol.5, No.5, 647-661, who describe the starch binding domain of *A.niger* glucoamylase.

Svensson et al., Biochem. J. (1989), 264, 309-311, describe the sequence homology between putative starch binding domains from α -amylase from *Streptomyces limosus*, β -amylase from *Clostridium thermosulfurogenes*, glucoamylase from *A.niger*, maltogenic α -amylase from *Bacillus stearothermophilus*, maltotetraose-forming amylase from *Pseudomonas stutzeri*, CGTase from *Bacillus*, CGTase from *Klebsiella pneumoniae* and glucoamylase from *Rhizopus oryzae* (the latter being a N-terminal starch binding region). It has also been suggested that some conserved tryptophan residues and the amino acids directly adjacent thereto may play an important role in

11-06-1999

EP99201862.2

DESC

3

starch binding, vide Goto et al., Appl. Environ. Microbiol., 1994, p.3926-3930; Williamson et al., Biochemistry 1997, 36, 7535-7539, and Chen et al., Protein Engineering, vol.8, 1049-1055 (1995).

WO 98/16190 describes fusion products of enzymes and one or more starch binding domains, as well as oral care compositions that contain such fusions. The fusions are prepared by expression of an appropriate expression vector in a suitable microorganism.

Chen et al., in Gene 99, (1991), 121-126, and in Biotechnol. Prog. 1991, 7, 225-229, describe a fusion of β -galactosidase and the starch binding domain from an *Aspergillus* glucoamylase, plasmids encoding such a fusion, and expression of said fusion in *E.coli*. The starch binding region is used to increase the affinity of β -galactosidase for starch granules, in particular as an affinity tail for recovery or enzymatic immobilisation using native starch granules as an absorbent.

Dalmia et al., in Biotechnology and bioengineering, Vol.47, pp.575-584 (1995), describe fusions of β -galactosidase and the starch binding domains of glucoamylase I of *Aspergillus awamori* and of cyclodextrin glucanotransferase (domain E of CGTase) from *Bacillus macerans*, respectively, plasmids encoding said fusions, and expression of said fusions in *E.coli*. The fusion proteins thus obtained are said to bind specifically to potato starch, corn starch, and cross-linked amylose. As a possible application, the use of the starch binding domains as an "affinity tag" is suggested. Similarly, Dalmia et al., in Enzyme Microb. Technol., 1994, vol.16, describe fusions containing a starch binding domain from *A.niger* glucoamylase, which is again used as an affinity tail to facilitate the one-step purification of the target β -galactosidase.

The use of cellulose binding domains as an affinity tag for protein purification (i.e. a fusion of a cellulose binding domain from a cellulase and α -galactosidase) has also been described in the art, vide Ong et al., Trends in Biotechnology, 7, 239-243 (1989).

However, when the above references describe fusions of a starch binding domain and an enzyme, said fusions are expressed in a micro-organism such as *E.coli*. Also, the starch binding domain is included only as a "tail" or "tag" in order to facilitate the isolation and purification of the desired enzyme activity from the bacterial culture medium.

The expression and use of such fusions in plants *in situ*, in particular in association with starch granules and/or plastids, is not mentioned in the art. In particular, the expression in plants of fusions containing enzymes that can alter the properties of the starch (granules) present in the plant (for instance for the *in vivo* production of improved starches) has not been described or suggested.

It is general object of the invention to express desired proteins or polypeptides in plants *in vivo* in association with starch produced or present in the plant, in particular in association with the starch granules produced or present in a plant.

A further object of the invention is to provide a method for obtaining transformed plants that produce modified starches.

Other objects of the invention will become clear from the description and figures hereinbelow.

The above objects are achieved by expressing a desired protein or polypeptide in the plant as a fusion with one or more starch binding domains.

In its broadest sense, the invention relates to a method for expressing a desired protein or polypeptide in a plant, in which the protein or polypeptide is expressed as a fusion with at least one starch binding domain.

In particular, the invention relates to a method for expressing a desired protein or polypeptide in a plant, such that the desired protein or polypeptide becomes associated and/or can be obtained in association with any starch produced by or present in said plant, and in particular with any starch granules produced by or present in said plant or any part of said plant, in which the protein or polypeptide is expressed *in vivo* in the plant or any part thereof as a fusion with one or more starch binding domains.

More specifically, this method comprises the steps of:

- a) providing a genetic construct comprising at least one nucleotide sequence encoding the desired protein or polypeptide combined with at least one nucleotide sequence encoding a starch binding domain;
- b) transforming a plant with said genetic construct;
- c) expressing said genetic construct in the plant *in vivo*.

Furthermore, as the starch granules in a plant will usually be present in cellular organelles within the plant cell, and in particular in plastids such as amyloplasts, chloroplasts and/or chromoplasts, the method of the invention can also be used to express a desired protein or polypeptide in association with such an organel.

11-06-1999

WAGENING

EP99201862.2

DESC

5

In a further aspect, the invention relates to a fusion of at least one desired protein or polypeptide and at least one starch binding domain, as expressed in - or more generally as present in - in a plant or in any part of a plant, including the seeds, leaves, roots (including tuberos roots), tubers, stems, stalks, fruits, grains or flowers, and in particular the honey-producing parts of flowers, of a plant.

A further aspect of the invention relates to a genetic construct suitable for transforming a plant, comprising at least one nucleotide sequence encoding a desired protein or polypeptide combined with at least one nucleotide sequence encoding a starch binding domain.

Yet other aspects of the invention comprise a plant that has been transformed with a genetic construct as mentioned above, or a descendant thereof, and/or a plant that expresses a fusion as described above, preferably in conjunction with any starch (granules) present in or produced by said plant.

Further aspects of the invention will become clear from the description hereinbelow.

The protein or polypeptide that is expressed via the method of the invention can be any polypeptide or protein known per se, and is preferably an enzyme. The protein or polypeptide may or may not by itself have natural affinity for starch or starch granules. If the protein or polypeptide essentially does not have any affinity for starch (granules), or has only a very low affinity, the method of invention may be used to provide the protein or polypeptide with such affinity. If the protein or polypeptide has some natural affinity for starch (granules), such as enzymes that naturally interact with starch or starch granules, the method of invention may be used to increase said affinity, and/or to alter said affinity. For instance, it is known that starch may occur in different forms or states, which may for instance differ in crystallinity (i.e. amorphous vs crystalline) and/or in one or more other properties such as glucan content or degree of branching, and such different forms of starch may also form different parts or regions of one and the same starch granule. The method of the invention may therefore also be used to alter (such as increase or decrease) the affinity of enzymes for one or more such specific forms of starch, both in absolute terms as well as in relative terms (i.e. compared to the affinity for other forms of starch). In this way, the affinity of the enzyme for different starch granules and/or for different parts or regions of a single starch granule may also be altered (again both in absolute or in relative terms).

11-06-1999

AGENINGEN

EP99201862.2

DESC

6

Therefore, a further aspect of the invention relates to a method for providing a protein or polypeptide with affinity for starch (granules), and/or for increasing the affinity of a protein or polypeptide for starch (granules), and/or for altering the affinity of a protein or polypeptide for starch (granules) comprising expressing the protein or polypeptide in a plant as a fusion with at least one starch binding domain.

More specifically, this aspect of the invention comprises the steps of:

- a) combining a nucleotide sequence encoding the protein or polypeptide with at least one nucleotide sequence encoding a starch binding domain, so as to provide a genetic construct encoding a fusion of the protein or polypeptide and the at least one starch binding domain;
- b) transforming a plant with said genetic construct;
- c) expressing said genetic construct in the plant.

The (gene(s) encoding) the desired protein or polypeptide may be derived from any source, including from plants, animals, fungi, algae, yeasts, bacteria and/or other micro-organisms, and may be homologous or heterologous to the plant in which the fusion is expressed.

Alternatively, (genes encoding) variants or mutants of such proteins or polypeptides may be used, such as those known per se in the art and/or obtainable via genetic manipulation techniques. These include mutant enzymes with altered properties compared to the enzyme from which they have been derived, such as altered substrate binding activity, altered substrate specificity, altered conversion properties and/or altered kinetic characteristics.

Furthermore, instead of the full protein or polypeptide as mentioned above, one or more fragments, parts, regions or domains thereof may be incorporated in and/or expressed as part of the fusions of the invention. Preferably, these fragments, parts, regions or domains part(s) are such that after expression, the fusion of the invention still provides the desired biological activity of the protein or polypeptide.

According to one preferred embodiment of the invention, the protein or polypeptide is an enzyme that can 'interact with' starch or starch granules, by which is meant that the enzyme can convert, modify, alter, degrade or otherwise influence the starch, the starch granule or the structure or interactions thereof.

Some non-limiting examples of such enzymes different kinds of amylases (i.e. alpha-, beta-, gluco-, iso-, etc.), CGTases, (neo)pullulanase, amylomaltase, glucan- and

11-06-1999

WAGENING

EP99201862.2

DESC

7

levansucrase, branching enzyme (such as glgB), phosphorylating enzymes such as certain kinases, oxidative enzymes (such as oxidases and dehydrogenase), other starch-decoloring enzymes (including those suitable for the transfer of sulfate and acetyl groups) and glycosyltransferases.

- 5 These enzymes may be obtained from any suitable source including bacteria, yeasts, fungi, algae and other micro-organisms including *Escherichia coli*, *Bacillus* (*subtilis*, *cereus*, *polymyxa*, *stearothermophilus*, *licheniformis*, *firmus/lentus*, *circulans*, *macerans*), *Aspergillus* (*niger*, *oryzae*, *kawachi*), *Klebsiella aerogenes*, *Streptomyces limosus*, *Pseudomonas* (*saccharophila*, *amyloderamosa*, *stutzeri*), *Clostridium* (*thermohydrosulfuricum*, *thermosulfurogenes*), *Microbacterium*, *Thermoanaerobacterium thermosulfurigenes*, *Saccharomyces cerevisiae*, *Rhizopus oryzae*, *Streptococcus mutans*, *Leuconostoc mesenteroides*, *Neisseria polysaccharea*, *Aureobasidium pullulans*.
- 10

- The enzymes may also be derived from higher forms of life including plants and animals. Some examples of plant-derived enzymes include starch synthases, starch-branching and -debranching enzymes (including isoforms thereof, such as BE-I and BE-II), disproportionating enzymes, putative potato kinase (sometimes also referred to as R1) etc., as well as the further enzymes mentioned in A.M. Smith et al. and C. Martin and A.M. Smith, above.
- 15

- 20 Other suitable starch-altering enzymes are for instance described in US-A-5,665,585.

- Some non-limiting examples of other proteins or polypeptides (i.e. non-enzymatic) that may be expressed in association with starch (granules) according to the invention include proteins or polypeptides derived from micro-organisms, plants or animals, such as receptors (such as estrogen receptors and in particular plant hormone receptors) and/or structural proteins, such as protein "zippers".
- 25

According to one embodiment of the invention, the desired protein or polypeptide is not β -galactosidase or another reporter enzyme.

- The term 'starch binding domain' or 'SBD' is well-known in the art, for instance from the references cited hereinabove, and is generally used to denote any part, region or domain of a protein or polypeptide, and in particular of an enzyme, that has natural affinity to (i.e. that binds to, attaches to, complexes with or otherwise associates with) starch or starch granules, or more generally with polymers of glucans.
- 30

Any naturally occurring starch binding domain, or any part or fragment thereof that still has affinity for starch (granules), may be used, as well as variants or mutants thereof.

As such, the starch binding domain(s) used in the invention may be derived from
5 any protein or polypeptide known per se that contains one or more starch binding domains, including proteins or polypeptides derived from plants, animals, fungi, algae, yeasts, bacteria and/or other micro-organisms. The starch binding domains may be homologous or heterologous to the plant in which the fusion is expressed. Preferably, a starch binding domain of an enzyme is used, more preferably an enzyme derived from a
10 bacterium, yeast, fungus or (other) micro-organism, or of a plant, such as GBSSI, which occurs in many different plants.

Some non-limiting examples of enzymes from which the starch binding domains may be derived are mentioned in the prior art indicated above, and may further include enzymes from bacteria, yeasts, fungi, or other micro-organisms such as the
15 cyclodextrin glycosyl transferases ("CGTases"), for instance from *Bacillus circulans*, *Aspergillus awamori*, *Aspergillus kawachi*, *Klebsiella pneumoniae* or *Bacillus stearothermophilus*; SBDs derived from thermostable enzymes, such as the CGTase of *Thermoanaerobacterium thermosulfurigenes*; glucoamylases, for instance from *Aspergillus niger* (which is reinforced by a disulfide bridge); glucoamylase from
20 *Rhizopus oryzae*, alpha-amylase from *Streptomyces limosus*, beta-amylase from *Clostridium thermosulfurogenes*, maltogenic alpha-amylase from *Bacillus stearothermophilus*, maltotetraose-forming amylase from *Pseudomonas stutzeri*; as well as enzymes from plants or animals such as granule-bound starch synthase I (GBSSI), partially granule-bound SSI, SSII, and SSIII, a putative kinase (R1), and
25 other granule-associated enzymes; as well as engineered forms of such domains.

Other suitable starch binding domains include for instance those derived from the enzymes mentioned in US-A-5,665,585, as well as the natural starch binding domains and variants thereof described by Penninga et al., Lawson et al., Sorimachi et al., Svensson et al., Goto et al., Williamson et al., and Chen et al., and Dalmia et al, above.

30 It is to be understood that although it is preferred in the invention to use only those parts or regions of the abovementioned enzymes, proteins or polypeptides that form the "starch binding domain(s)", it is not excluded that one or more other parts or sequences of the original enzyme may also remain or be present in the fusions of the

11-06-1999

WAGENING

EP99201862.2

DESC

9

invention (i.e. attached to the starch binding domain). It may even be possible to use the full amino acid sequence of the original enzyme as a starch binding domain in the invention, i.e. as a fusion with the desired enzyme, protein or polypeptide. When such further parts or sequences are (still) present, these most preferably (no longer) show any biological activity per se. When a full sequence is used as a starch binding domain, this sequence has preferably been made devoid of biological activity (e.g. made catalytically inactive), although again the invention in its widest sense is not limited thereto.

Some preferred examples of enzymes that can be used as starch binding domains in the invention (i.e after they have been made catalytically inactive) include catalytically inactive GBSS I or gbSSII.

Generally, such starch binding domains are polypeptides of about 95 to about 105 amino acids amino acids, although their size is not essential in the invention. For instance, when catalytically active enzymes such as GBSSI or gbSSII or parts thereof are used, they may have a size of about 525-770 amino acids. For some specific applications, such large strach binding groups may be preferred, but generally smaller size SBD's (i.e. up to about 150 a.a.) will be preferred.

According to one embodiment, the starch binding domain used in the invention contains at least one of the minimal sequences shown in Figures 1 or a variant or mutant thereof, for instance in which one of the tyrosine residues in binding site 2 have been replaced by tryptophan.

Other suitable starch binding domains can be identified by means of sequence alignment of the above minimal sequences of Figures 1A with a known database, for instance an alignment program known per se such as BLAST or PC gene. In general, any domain or region of an enzyme that contains a sequence that has a sequence homology of more than 50%, preferably more than 70%, more preferably more than 90% with the abovementioned minimal sequence as can be used (in which a deletion or insertion is counted as a single mutation).

Preferably, any binding domain used in the invention has an affinity for starch (granules) (expressed as K_{ad} -value) of more than 10, preferably more than 15 ml/g, as measured by means of adsorption isotherm methodology as described by Chen et al, for instance in the Protein Engineering-, Biotechnol. Prog.- and/or Gene-references mentioned above. In general, this involves measuring the adsorption of the binding

11-06-1999

GENINGEN

EP99201862.2

DESC

10

domain to native starch at different concentrations of the protein (i.e. of the binding domain). After mixing the protein (for instance in concentrations ranging from 0,1 to 1,0 mg/ml) and the starch (for instance 0,1g) in a suitable aqueous medium (for instance 1 ml total volume), the mixture is shaken for a suitable period of time (for instance 20 min to 1 hr) at a suitable temperature (for instance 4°), after which the mixture is subjected to centrifugation (for instance 17400 g for 20 min), after which the protein concentration of the supernatant is assayed and the amount of adsorbed protein is determined by subtraction. Values for the adsorption constant (K_{ad}) can then be derived from the slopes of the linear adsorption isotherms, optionally by comparison with a reference protein.

Also, preferably, the entire fusion of the invention also has an affinity for starch (again expressed in terms of K_{ad} - value) of more than 10, preferably more than 15 mL/g.

With respect to the affinity for starch and the K_{ad} -value, it will further be clear to the skilled person that these may also be influenced by factors such as the type of starch used, the presence of glycosylation, the size of the fusion, the type of linker (if any) etc.. Also, other assays for determining the affinity of a protein to starch have been described in the art, and these can also be used to identify suitable domains or regions.

In another embodiment of the invention, and as an alternative for starch binding domains, domains or regions with affinity for other glucan polymers such as cellulose, amylose, or amylopectin, glycogen, mutan, dextran, nigeran, pullulan, or affinity for fructan, pectin, xylan, or (mixed linkage) beta-glucan polymers may be used. These include for instance enzymes or proteins for glucan polymers other than starch such as the glucan binding domains of sucrases, cellulose binding domains (which however have no or very low affinity for starch, and are therefore less preferred), granule-bound starch synthase I or part thereof, as well as engineered sucrose porin, maltose binding protein, maltoporin and/or lamB from *E.coli*, which have affinity for maltose/maltodextrin (The latter, however, have the disadvantage of a large size, i.e. about 400-600 amino acids, and also have little or no affinity for so-called raw starch).

Such cellulose binding domains, maltose binding proteins etc. can be used/expressed in a fusion of the invention analogously as described herein for the starch binding domains, to provide fusions with affinity for, and/or that can associate with, cellulose and maltose/maltodextrin, respectively. However, the use of starch

11-06-1999

WAGENING

EP99201862.2

DESC

11

binding domains, so that the resulting fusion can associate with starch (granules), is much preferred.

In the fusions expressed according to the invention, the desired protein or polypeptide may be fused directly with the one or more starch binding domains, or via a linker sequence, i.e. a sequence of 1-100, preferably 4-60 (optionally glycosylated) amino acids that connects the protein or polypeptide with the one or more starch binding domains. The linker sequence may also act as a "hinge" and/or spacer, for instance in order to ensure that the presence of the starch binding domain or the binding thereof to the starch (granule) does not interfere with or detract from the desired activity or properties of the enzyme, protein or polypeptide (and visa versa).

In principle, any natural or synthetic amino acid sequence can be used as a linker, preferably a sequence that essentially does not interfere with either the affinity of the starch binding domain for the starch (granules), or with the desired activity of the enzyme.

Some non-limiting examples of suitable linker sequences include:

- the 40 AA linker sequence used in Trichoderma, comprising a flexible part and glycosylated rigid part;
- naturally occurring or synthetic sequences of alternating Pro-Thr;
- linker sequences containing major amount glycine residues or of (optionally O-glycosylated) serine or threonine residues.

Other suitable linkers are for instance described by P. Argos, in J. Mol. Biol. (1990), 211, p. 943-958.

The protein or polypeptide may also be bound to the one or more starch binding domains via an amino acid sequence that can be cleaved, i.e. chemically or preferably enzymatically. Examples are amino acid sequences that provide enzymatic cleavage site for enzymes such as thrombin, factor Xa, and collagenase can be mentioned. Such a sequence may also form part of a larger linker sequence as described above.

In the fusions of the invention, the one or more starch binding domains may be at the N'-terminus of the fusion, at the C' terminus of the fusion, or - if two or more starch binding domain are present - both. If a starch binding domain is used that is naturally (i.e. in the enzyme from which it has been derived) is at the C' terminus (or N' terminus), it is preferably also at the C' terminus (or N' terminus, respectively) of the fusion of the invention, although the invention is not limited thereto.

11-06-1999

INGENINGEN

EP99201862.2

DESC

12

Also, when a fusion to the invention contains two or more starch binding domains, these may be the same or different (such as different domains from the same enzyme or domains from different enzymes/sources) and may -if present at the same terminus - also be fused directly or via a linker sequence, that again may also act as a hinge and/or spacer.

Furthermore, although usually not preferred, it is also possible that a fusion of the invention (a "bifunctional fusion") contains two or more of desired enzymes, proteins or polypeptides, which may be fused directly, via a linker sequence, or via a sequence encoding one or more starch binding domains (besides any further starch binding domains that may be present at the 3' and/or 5' end). For instance, there are cellulases and xylanases with "internal" cellulase binding domains, which may be included "internally" between two catalytic centers of the bifunctional fusion.

The fusions of the invention are obtained by expression in a plant *in vivo* of a nucleotide sequence (genetic construct) that codes for the fusion of the invention. For this purpose, the plant may be or may have been transformed with said genetic construct, or may be a descendant (such as obtained via sexual or asexual multiplication, including crossing and/or other breeding techniques) of a plant that has been transformed with such a genetic construct, and that has inherited the genetic construct.

The plant may be any monocotylous or dicotylous plant in which the fusions can be expressed, but is preferably a plant that naturally contains or produces starch, and more preferably a plant that contains or produces starch granules, either throughout the entire plant or in any part thereof, including seeds, leaves, roots (including tuberos roots), tubers, stems, stalks, fruits, grains or flowers, and in particular the honey-producing parts of flowers; and such a plant is referred to herein as a 'starch granule producing plant'. As mentioned above, said starch granules will usually be associated with or present in specific organelles of the plant cell, and in particular the plastids, such as chloroplasts, amyloplasts and/or chromoplasts.

Some preferred non-limiting examples of starch granule producing plants suitable for use in the invention include economically important crops such as potato, sweet potato, cassava, pea, taro, sago, yam, banana and/or cereals such as rice, maize, wheat and barley; of which potato, sweet potato, maize and wheat are especially preferred.

11-06-1999

WAGENI

EP99201862.2

DESC

13

The starch granule producing plant may also be a plant, and in particular a genetically modified plant, that as such already produces a modified starch, such as transformed potato plants producing mutant amylose free ("amf") starch. For some applications, the use of a plant that produces starch granules that contain pores may be useful, in particular when the further use of the starch granules associated with the fusions of the invention involved diffusion of compounds in and out of the granules. For instance, some cereal starches are known to have natural pores. Also, the use of small granules, again as for instance in cereal starches, may also be advantageous with respect to diffusion of substances into the starch granule. Alternatively, a (pre)treatment with an enzyme that make pores or holes in starch granules, such as those of microbial origin, may be used.

A genetic construct encoding a fusion of the invention may be obtained by 'combining' the nucleotide sequence(s) encoding the at least one desired protein or polypeptide with at least one nucleotide sequence that codes for a starch binding domain, optionally with or via one or more sequences that encode a linker sequence as described above, in such a way that expression of the combined sequences in the desired plant leads to the formation of the fusion.

Generally, this involves covalently binding the nucleotide sequences in the same reading frame and in the same orientation, and in the correct order from the 5' end to the 3' end. This can be carried out using genetic manipulation techniques known per se, such as those described in Sambrook et al, "Molecular Cloning: A Laboratory Manual" (2nd.ed.), Vols. 1-3, Cold Spring Harbor Laboratory (1989); or F. Ausubel et al, eds., "Current protocols in molecular biology", Green Publishing and Wiley Interscience, New York (1987)

The one or more sequence(s) encoding the starch binding domains can be provided synthetically using known DNA synthesis techniques, but are preferably isolated from the organism from which the starch binding domain has been derived (i.e. in which it naturally occurs). Similarly, the sequence encoding the protein or polypeptide is also preferably isolated from a suitable biological source, and as such may be a nucleotide sequence encoding the mature protein, or a nucleotide sequence encoding a precursor thereof, that can be converted into the mature protein or polypeptide by post-translational modification(s) in the plant (i.e. as part of the encoded fusion).

11-06-1999

AGENINGEN

EP99201862.2

DESC

14

The genetic construct encoding the fusions of the invention may further contain all other elements known per se for nucleic acid sequences or genetic constructs, such as promoters or other control elements, terminators, translation or transcription enhancers, integration factors, signal sequences, selection markers, etc., that are preferably suited for use in (the transformation of) the host plant. The sequences that encode these further elements of the construct may again be either isolated from a suitable biological source, or provided synthetically. Examples of suitable elements are for instance described in DE-A-195 34 759, WO 91/19808, US-A-5,349,123, US-A-5,750,875 and WO 92/14827.

10 The one or more nucleotide sequences encoding the further elements of the construct can again be combined with the nucleotide sequence encoding the fusion in a manner known per se, such as described in Sambrook et al., Ausubel et al., DE-A-195 34 759, WO 91/19808, US-A-5,349,123 or US-A-5,750,875.

15 Preferably, they are combined in such a way that - after transformation - the construct can be used for the expression of the fusion in the desired plant. Generally, this involves combining the control elements and any further elements with the sequence encoding the fusion in an operable manner, i.e. in the same reading frame and in the same orientation, and in the correct order from the 5' end to the 3' end.

20 The promoter can be any promoter that is able to control/induce the expression of the fusion in the intended plant, including constitutive and inducible promoters, and may be homologous or heterologous to said plant.

25 Also, a promoter may be used that directs the expression of the fusion to a specific part or tissue of the plant, and in particular to a tissue or part of the plant where starch (granules) are formed or present, including the including seeds, leaves, roots (including tuberos roots), tubers, stems, stalks, fruits, grains or flowers. and in particular the honey-producing parts of flowers, etc.. Furthermore, a promoter may be used that induces expression of the fusion during a specific period in the life cycle of the plant. For instance, in potato, a promoter may be used that specifically directs the expression of the fusion in or to the tuber, and/or that allows for expression only during
30 the time the plant forms its tubers.

Examples of suitable promoters include the CaMV promoter, GBSS promoter, patatin promoter, Ubiquitin promoter, ST1 promoter, TR1 promoter, napin promoter, as well as for instance the promoters described in DE-A-195 34 759, WO 91/19808,

11-06-1999

WAGENIN

EP99201862.2

DESC

15

US-A-5,349,123, US-A-5,750,875 and WO 92/14827. For specific expression of foreign genes in potato tubers, reference is made to for instance EP 0 375 092 and Rocha-Sosa et al., EMBO J. 8, 23-29 (1989).

The construct of the invention may also comprise one or more sequences that
5 encode signal proteins, including pre-, pro- of prepro-sequences. These usually precede the sequence encoding the fusion, such that the fusion is expressed as a (further) fusion with these signal proteins. The signal sequence may ensure any post-translational modifications required for the formation of the mature fusion (i.e. of the
10 protein/polypeptide and/or the starch binding domain(s) part thereof), and/or may specifically direct the expressed fusion to a desired part or organel within the plant or plant cell, and in particular to the starch granule(s). In particular, signal sequences for plastide targeting, such as for amyloplast, chloroplast or chromoplast targeting can be used, or signal sequences for targeting the vacuole. Some non-limiting examples thereof include the small subunit RuBisCo, GBSS transit peptides and sporamine
15 transit peptide.

According to one preferred embodiment, the genetic construct encoding the fusion is preferably in a form suitable for transformation of a plant, such as a vector or plasmid. As such, the construct is preferably such that upon transformation it is incorporated into the (genomic) DNA of the plant. However, the construct may also be
20 in any other form that can provide for expression of the fusion in the plant, and that preferably also can be stably and/or independantly maintained and/or replicated in the plant, and/or inherited from one generation of the plant to the next. The construct is preferably further in a form that can be stably and/or independantly maintained and/or replicated in any organism to be used for constructing or selecting the construct and/or
25 to be used in transforming the plant, such as *Agrobacterium*.

A further aspect of the invention therefore relates to a bacterium, virus or other organism suitable for transforming a plant, containing a genetic construct as defined above, and preferably capable of transferring said construct into a plant. The organism may for instance be a strain of *Agrobacterium*.

30 After construction, the construct is transformed into the desired plant, preferably a starch granule producing plant as defined above. Any technique for the transformation of a plant known per se can be used. Examples thereof include transformation using *A. tumefaciens* or *A. rhizogenes*, electroporation of tissues and/or

11-06-1999

AGENINGEN

EP99201862.2

DESC

16

protoplasts, particle bombardment, use of virusses for DNA delivery, etc., as well as the techniques described in DE-A-195 34 759, WO 91/19808, US-A-5,349,123, US-A-5,750,875 and WO 92/14827

After transformation, a plant is (re)generated from the transformed cells or tissue and the construct is expressed in the plant or part thereof, optionally upon induction thereof in a suitable manner.

The invention therefore also relates to a method for providing a plant that expresses a fusion as described above, comprising at least one step of:

- a) transforming a plant with a genetic construct as described above, such that said genetic construct is expressed in the plant or at least part thereof; and optionally further comprising at least one step of:
- b) providing descendants and/or further generations of the thus transformed plant, for instance via sexual or asexual multiplication, including crossing and/or other breeding techniques.

The invention also relates to seeds, tubers, seedlings, stakes (e.g. for cassava) or other cultivating material of such a transformed plant.

Upon expression, the fusion will usually become 'associated with' any starch granules present in the plant, by which is meant that the fusion attaches to, binds to, complexes with or otherwise combines with the starch granule, i.e. via the one or more starch binding domains present in the fusion. Said association may be such that the fusion is present on the surface of the starch granule, and/or incorporated into (such as by encapsulation or enclosure) the starch granule (e.g. during the biosynthesis thereof). The association(s) thus obtained will be collectively referred to hereinbelow as the 'complex'. One preferred embodiment of the invention concerns these complexes, methods for their preparation by expression in a plant, and plants that express these complexes.

As mentioned above, the starch granules in a plant will usually be present in or associated with specific organelles within the plant cell, and in particular the plastids, such as the amyloplasts, chloroplasts or chromoplasts. It should be understood that when in the present description and claims mentioned is made of a "starch granule", this is also meant to include (the starch granules as present in) these organelles. The term "complex" as mentioned herein therefore also includes complexes of fusions of the invention and such organelles.

11-06-1999

WAGENIN

EP99201862.2

DESC

17

Therefore, in another aspect, the invention relates to a method for producing a complex of at least one protein or polypeptide and a starch granule, comprising at least one step of:

- 5 a) expressing the protein or polypeptide as a fusion with at least one starch binding domain, in a plant that contains or forms starch granules; and optionally comprising at least one further step of:
- b) isolating the protein or polypeptide from the plant or any part thereof as a complex of the fusion and the starch granule.

10 A further aspect of the invention relates to a complex, comprising a fusion of a protein or polypeptide and at least one starch binding domain, associated with a starch granule.

In particular, this aspect of the invention relates to such a complex as expressed in/obtained from a plant via the method described above.

15 Yet another aspect of the invention therefore relates to a method for providing a plant that can produce a complex of a fusion as described above and a starch granule, comprising at least one step of:

- a) transforming a starch granule producing plant with a genetic construct as described above, such that said genetic construct is expressed in the plant or at least part thereof;
- 20 and optionally further comprising at least one step of:
- b) providing descendants and/or further generations of the thus transformed plant, for instance via sexual or asexual multiplication, including crossing and/or other breeding techniques.

25 The invention further relates to the complex-producing transformed plants thus obtained, or any descendant thereof, as well as cultivation material of said plant, including seed, tubers, stakes or seedlings.

Usually, and preferably, the fusion and the starch granule will already associate *in vivo*, so that they can be obtained/isolated together, using techniques known per se for the isolation of the starch granules from the plant or plant material. However, the invention is not limited thereto. For instance, the complex may also be formed during or as a result of the isolation/further processing of the plant or plant material, for instance when the fusion is expressed in a part of the plant (cell) separate(d) from the starch granule. In yet another embodiment, starch granules, optionally combined with a

11-06-1999

AGENINGEN

EP99201862.2

DESC

18

further carrier or matrix, can be used to selectively isolate the fusion from the plant (material), i.e. by a method comparable to an affinity technique.

Again, instead of isolating the starch granules, it may be easier and/or advantageous to isolate the organelles in which the starch granules are present, again in a manner known per se, so as to provide complexes of the starch granule containing
5 a manner known per se, so as to provide complexes of the starch granule containing organel and the fusion of the invention. These may also be used as such in any subsequent application(s).

The above aspect of the invention is particularly suited for producing a desired protein or polypeptide in for instance potato tuber, cassave root, sweet potato tuber,
10 grains of maize, wheat and barley, as well as in peas, etc.. After the tubers, roots or grains produced by the plants expressing the fusions have been harvested, the protein or polypeptide can be conveniently isolated therefrom as a complex with the starch granules. This aspect thus provides for a very efficient production and isolation of any desired protein or polypeptide, and may be used to produce such a polypeptide or
15 protein in major and/or commercial amounts.

The complex thus obtained may be processed further, for instance for further purification, in which the fact that (essentially only) the desired protein or polypeptide is present as a complex with the starch granules may be used with advantage. The fusion may also be separated from the starch granules, and/or the fusion may be
20 cleaved (i.e. at a suitably situated enzymatic cleavage site as described above) in order to provide the desired protein or polypeptide.

However, it may also be advantageous to use the complex as such, for instance when the polypeptide is an enzyme. The starch granule or organelle can then act as a matrix or carrier for the desired enzyme activity, which may for instance facilitate the
25 handling or use thereof in any subsequent application, provide for an easily reusable enzyme preparation, or improve the half life of the enzyme.

Compared to the expression of fusions in bacterial expression hosts such the *E.coli* strains described in the art, expression of the desired proteins or polypeptides according to the invention in (edible) plant material such as potato tubers may be
30 advantageous from a safety standpoint. Also, expression of the desired protein or polypeptide in a plant host compared to a bacterial host may be advantageous for some applications, for instance when a bacterial host does not carry out all desired post-translational modifications or conversly degrades or otherwise detracts from the desired

11-06-1999

WAGENI

EP99201862.2

DESC

19

protein or polypeptide. For such applications, the method of the invention may provide a valuable alternative.

According to another preferred embodiment of the invention, the plant used to express the fusions of the invention is a starch producing plant, in particular a starch granule producing plant, and the protein or polypeptide expressed as part of said fusion according to the invention is an enzyme that can interact with starch or starch granules.

In this embodiment, after expression, the fusions preferably associate (either *in vivo* and/or during processing of the plant or plant material) with the starch (granules) via the one or more starch binding domains, after which the enzyme can interact with the starch (granule), for instance to convert, modify, alter, degrade or otherwise influence the starch, the starch granule or the (primary) structure or interactions thereof, resulting in (a plant or plant material that can be used to provide) a modified starch, i.e. a starch different from the starch naturally provided by the plant in at least one property thereof, such as crystallinity, branching degree, protein content, glucan composition, oxidation, phosphorylation, etc..

In particular, in this aspect of the invention, the enzyme that can interact with the starch (granules) is an enzyme that does not occur naturally in (i.e. that is heterologous to) the original starch producing plant, but is for instance an enzyme derived from another plant or from a bacterium, fungus or (other) micro-organism as described above.

In this way, the invention can be used to provide a transformed plant that *in vivo* produces starches that are modified or altered (i.e. compared by the starch naturally produced by the original starch producing plant).

Therefore, yet another aspect of the invention relates to a method for providing a plant that contains or produces a modified starch and/or modified starch granules, comprising at least one step of:

- a) transforming a starch producing plant, in particular a starch granules producing plant, with a genetic construct comprising at least one nucleotide sequence encoding an enzyme that can interact with starch and/or starch granules and at least one nucleotide sequence encoding a starch binding domain, such that said genetic construct is expressed in the plant or at least part thereof; and optionally further comprising at least one step of:

11-06-1999

AGENINGEN

EP99201862.2

DESC

20

- b) providing descendants and/or further generations of the thus transformed plant, for instance via sexual or asexual multiplication, including crossing and/or other breeding techniques.

5 The invention further relates to the transformed plant producing modified starch (granules) thus obtained, or any descendant thereof, as well as cultivation material of said plant, including seed, tubers, stakes or seedlings.

This aspect of the invention also comprises a method for producing a modified starch and/or modified starch granules, comprising at least one step of:

- 10 a) cultivating a transformed plant that produces a modified starch and/or modified starch granules as described above, or a descendant thereof:
and optionally further comprises at least one step of:
b) isolating the modified starch or starch granules from the transformed plant or from any part thereof, such as its seeds, leaves, roots (including tuberos roots), tubers, stems, stalks, fruits, grains or flowers.

15 This aspect of the invention can in particular be used to provide potato, sweet potato, cassava and/or cereals such as maize, rice, wheat, and barley or other economically important crops that produce modified starches, i.e. in their tubers or seed, and such tubers or seeds containing modified starches form a further aspect of the invention. Another aspect of the invention resides in the modified starches thus
20 obtained.

Alternatively, the invention may be used to provide starch granules that already contain one or more enzymes that can interact with starch, or plant material that contains such starch granules, i.e. attached to or incorporated within the starch granule via the one or more starch binding domains. After harvesting of the plant, plant
25 material and/or starch granules, the enzymes may then be used *in vitro* to alter the properties of the starch (granule), to provide a modified starch.

For some applications, this embodiment may offer advantages over the *in planta* production of modified starches, for instance when specific conditions (such as temperature, pH optimum, the presence of certain co-factors, etc) are required for the
30 starch converting activity of the enzyme, and/or when greater control over the enzymatic starch conversion(s) is desired. For instance, enzymes such as thermostable α -amylases and isoamylase may have an optimal temperature range for conversion that

11-06-1999

WAGENI

EP99201862.2

DESC

21

may not be achieved *in planta* (i.e. in the field); a suitable temperature can then be applied during subsequent processing of the harvested plant material.

In a sense, this embodiment combines features of the two aspects of the invention described above, in that on the one hand the invention is used to
5 express/provide a "complex" of a starch (granule) and a fusion of a starch-converting enzyme and one or more starch binding domains; whereas on the other hand the enzymatic activity present in the complex thus obtained is then used – through post-harvesting modification(s) - to provide a modified starch.

This aspect of the invention therefore relates to a method for producing a modified
10 starch and/or modified starch granules, comprising at least one step of:

- a) cultivating a transformed plant that expresses a fusion of at least one starch binding domain and at least one enzyme that can interact with starch (granules), in which the starch (granule) and the fusion preferably form a complex;
- b) harvesting the plant and/or any part of said plant that contains starch (granules) and
15 the fusion, such as the seeds, leaves, roots (including tuberos roots), tubers, stems, stalks, fruits, grains or flowers (in particular the honey-producing parts thereof);
- c) subjecting the plant or plant material, or any fraction or preparation obtained therefrom that contains the starch (granule) and the fusion, to conditions such that the enzyme can interact with the starch (granule) to provide modified starch or
20 starch granules;

and optionally comprises the further step of :

- d) isolating the modified starch or starch granules thus obtained.

In this embodiment, the "post-harvest" modification of the starch of step c) can for instance be carried as part of the isolation of the starch (granules) from the plant or
25 plant material, or can be carried out in a separate processing step on either the harvested plant or plant material, on any fraction or preparation obtained therefrom that contains both the starch (granule), or on the isolated starch (granule), i.e. while it is still associated with the fusion (i.e. as a complex as described herein).

The invention further relates to the modified starch (granules) thus obtained.

30 Furthermore, and analogous to the above described interaction with the starch (granules), the enzymatic activity expressed as part of the fusion may also convert, modify, alter, degrade or otherwise influence any other compound(s) present within the plant, or any other part(s) or biological function of the plant. The enzymatic activity

11-06-1999

AGENINGEN

EP99201862.2

DESC

may also provide for the *in vivo* synthesis of one or more desired compounds within the plant, which may or may not occur naturally in the original plant.

Therefore, in its broadest sense, this embodiment of the invention may be used to provide the plant with (i.e. to express within the plant) any desired enzymatic and/or biosynthetic activity, in association with any starch (granules) present within the plant.

According to yet another embodiment of the invention, only a nucleotide sequence encoding one or more starch binding domains as described above is expressed in a plant. According to this embodiment, the starch binding domain, which usually will be heterologous to the plant, upon expression preferably associates with any starch (granules) present in the plant, so as to make the starch granules less accessible to or a less favorable substrate or one or more starch converting enzymes that are naturally present in the plant. More generally, the starch binding domains thus expressed may compete with the native starch converting enzymes present in the plant (i.e. for "space" on or in the starch granules), thus influencing the native starch conversion processes.

In this way, the method of the invention can be used to inhibit, alter, modify or otherwise influence one or more of the native biosynthetic pathways in the plant involved in the starch (granule) biosynthesis or metabolism, which again can result in a (plant producing a) modified starch. Also, the interaction of the starch binding domain(s) and the starch (granule) can again lead to an alteration or modification of the starch granule, such as changes in the (primary) structure, in the amylose content, and/or in crystallinity etc..

This aspect of the invention therefore generally comprises transforming a plant with a nucleotide sequence that codes for at least one starch binding domain, such that said starch binding domain is expressed in the plant, and preferably associated *in vivo* with any starch (granules) present in the plant or in any part thereof.

The nucleotide sequence is preferably in the form of a genetic construct as described above (but containing only the one or more sequences encoding a starch binding domain, optionally linked via a linker sequence, and any further elements known per se for such constructs as described above). Also, as more generally mentioned above, the 'starch binding domain' used in the may also include parts of the enzyme, protein or polypeptide from which the domain was derived, or may even be a full protein that has been made catalytically inactive, such as catalytically inactive GBSS I or the partially granule bound SSII (also sometimes referred to as gbSSII).

11-06-1999

WAGENINGEN

EP99201862.2

DESC

Therefore, yet another aspect of the invention relates to a method for providing a plant that contains or produces a modified starch and/or modified starch granules, comprising at least one step of:

- 5 a) transforming a starch producing plant, in particular a starch granules producing plant, with a genetic construct comprising at least one nucleotide sequence encoding a starch binding domain, such that said genetic construct is expressed in the plant or at least part thereof;
- and optionally further comprising at least one step of:
- 10 b) providing descendants and/or further generations of the thus transformed plant, for instance via sexual or asexual multiplication, including crossing and/or other breeding techniques.

The invention further relates to the transformed plant producing modified starch (granules) thus obtained, or any descendant thereof, as well as cultivation material of said plant, including seed, tubers or seedlings.

- 15 This aspect of the invention also comprises a method for producing a modified starch and/or modified starch granules, comprising at least one step of:
- a) cultivating a transformed plant that produces a modified starch and/or modified starch granules as described above, or a descendant thereof;
- and optionally further comprises at least one step of:
- 20 b) isolating the modified starch or starch granules from the transformed plant or from any part thereof, such as its seeds, leaves, roots, tubers, fruits, etc..

Again, this aspect can in particular be used to provide provide potato, cassava, sweet potato, taro, sago, yam, banana and/or cereals such as rice maize, wheat, barley or other economically important crops that produce modified starches, i.e. in their
25 tubers or seed, and such tubers or seeds containing modified starches, from a further aspect of the invention. Another aspect of the invention resides in the modified starches thus obtained, optionally in the form of a complex with one or more starch binding domains as obtained through expression of the above genetic construct.

Some non-limiting, preferred practical applications of the invention include:

- 30 - Production of amylose-free starch, in particular amylose-free potato starch.
- Introduction of starch-binding domains, or any other protein binding to starch granules, may be used as an alternative for antisensing starch-converting enzymes, such as the GBSS I gene (or for mutating the GBSS I gene), to obtain amylose-free

potato starch. In the *amf* mutant and the antisense GBSS I potato plants down-regulation of the amylose content occurs at the DNA and RNA level, respectively. In the SBD-expressing potato plants, reduction of the amylose content is based on competition between the expressed SBDs and GBSS I.

5 The inhibition of the starch-converting enzymes may be further manipulated (i.e. increased) by using tandem SBDs (instead of a single SBD) separated by an appropriate linker peptide. Depending on the affinity for raw starch of the SBD in comparison with GBSS I, a single SBD or a series of more than two SBDs may achieve an amylose-free potato starch.

10 - Modification of starch fine structure *in planta*.

Modification of the fine structure of starch can be achieved *in planta* by concentrating certain enzymes, equipped with an SBD, in the granule, or at the granule surface. Thus, next to targeting polypeptides to the amyloplast, also targeting within the amyloplast can be achieved. For instance, the use of

15 *Escherichia coli* glgB or a potato kinase (sometimes referred to as R1) fused with an SBD can increase the degree of branching and phosphorylation of the starch to a larger extent than the non-engineered proteins.

Generating (new) transgenic starches, which either may have different functional properties compared to the existing ones (such as increased freeze-thaw stability or altered rheological properties), or which may be regarded as a better precursor for derivatization processes than the WT or *amf* potato starch.

20

- Starch modification *in vitro*.

In addition to modification *in planta*, (further) modification of the fine structure of the starch may take place during or after processing of the harvested plant

25 (material). By introducing a protein in or on the surface of the starch granule, a complex of the invention that serves as a kind of 'precursor starch' can be obtained, which can be converted to the starch of interest after extraction of the starch from the plant (material). Thus, in this embodiment, the actual (or full) modification of the starch may not or may only in part be achieved *in planta*. The advantage of this is that (further) modification of the starch can take place under more controlled, and/or more extreme conditions than encountered in the plant *in vivo*, and/or in the presence of compounds such as reactants or co-factors which are not present in the amyloplast.

30

11-06-1999

WAGENIN

EP99201862.2

DESC

25

As an example thereof, the introduction of an oxidase/dehydrogenase as a fusion with one or more starch binding domains can be mentioned, yielding a more reactive starch after appropriate incubation *in vitro*. In this way, the derivatized starch can be obtained in a much more environmentally friendly manner than when chemical methods were applied.

5

- Starch granules with immobilized enzymes.

In this application, a foreign enzyme is introduced in or onto the starch granule, with the objective of using the granules as a carrier of enzymes. An enzyme of choice is fused to a SBD (with linker), and this fusion protein is incorporated into the starch granule during starch biosynthesis. Subsequently, these granules can be used to catalyze certain *in vitro* conversions in which reactants and products diffuse in and out of the granule. Such a procedure allows a simple separation of enzyme and products. In addition the enzyme can be re-used.

10

- Specialty support for affinity chromatography.

In this application, a protein without a catalytic activity is introduced in or onto the starch granule during starch biosynthesis. The SBD can be fused to any kind of receptor, and the forth-coming starch can subsequently be used as a support for affinity chromatography, i.e. small molecules can be specifically absorbed from complex mixtures such as culture filtrates or plant extracts, and eluted after several wash steps.

20

- Molecular pharming of industrial enzymes.

In this application the starch granule also serves as an affinity support, but in a different context. Industrially relevant enzymes are fused to a SBD (with linker), and can be produced in, for instance, the vacuole of plant cells. Upon disruption of the tissue, the enzymes are released from this compartment, and are able to contact the surface of granules released from the amyloplast. After washing, the fusion proteins may be eluted from the granules with a maltodextrin solution. It is also possible to release the enzyme (without SBD) with a highly specific protease.

25

- Production of cross-linked starches or starch granules.

In this application, protein-carbohydrate interactions (SBD-starch) are combined with protein-protein interactions (for instance, a leucine 'zipper').

30

Two possible uses thereof are for instance in reinforcing the granule and/or in creating a stronger starch gel after the granules are gelatinized.

11-06-1999

WAGENINGEN

EP99201862.2

DESC

26

For reinforcing the granule, a leucine zipper domain may be flanked by one or two SBDs. The zipper domains can create a second network in the granule, which is connected to the first (starch) by the SBD(s). Possibly, the melting temperature of the granule can be changed in this way. Reinforcing the starch structure in this way may also be useful for providing complexes of the invention that can for example be used in high temperature conversions. (Mutant starches like maize amylose-extender starch may also be useful in this respect)

For creating a stronger starch gel after gelatinization, a zipper domain may be flanked by a SBD on one side and a Maltose Binding Protein or 'MBP' on the other. The zipper domain is anchored in the granule by a SBD. After gelatinization, the zipper domains are connected, and this new network is attached to the starch network by MBPs. Gels with new properties may be realized in this way. In this aspect, the thermostability of the MBP may be of importance.

The invention will now be further illustrated by means of the Experimental Part given hereinbelow, as well as the Figures, in which:

- Figure 1 shows the amino acid sequences for some minimal sequences for starch binding domains suitable for use in the invention;
- Figures 2A-2E show some examples of genetic constructs according to the invention.

Experimental Part

A. Introduction

Starch is an important storage material in many plants, such as potato, sweet potato, cassava, pea, yam, taro, sago, banana and cereals such as maize, rice, and potato. It is deposited as crystalline granules which generally consist of two polysaccharides, amylose (generally 20-30%) and amylopectin (70-80%). Amylose is an essentially linear molecule which is composed of (1→4)-linked α -D-glucopyranosyl (α -D-Glcp) residues. Amylopectin is a highly branched molecule composed of a collection of α -(1→4)-glucan chains which are connected by α -(1→6)-linkages (the branch points). It is believed that branching occurs at regular intervals, in such a way that clusters of sidechains are formed. The sidechains can interact laterally with each other which is presumably the basis for the crystalline nature of the granule.

11-06-1999

WAGENING

EP99201862.2

DESC

- Current models explain granule growth by addition of single Glc residues to the non-reducing ends of a nascent amylopectin molecule, a reaction catalyzed by synthases. Once the glucan chains have reached a certain length, they are thought to be rearranged with the help of a number of enzymes such as branching enzyme (BE),
5 debranching enzyme (R-enzyme), and presumably also disproportionating enzyme (D-enzyme). As far as is currently known, starch-producing plant species can possess several isoforms of starch synthases (GBSS I, GBSS Ib, gbSS II, SSS I, and SSS III; GB=granule-bound, gb=partially granule-bound, S=soluble) and BE (BE I, BE IIa, and BE IIb).
- 10 The major differences between the potato starch synthase isoforms are the length of the N-terminal extension, and the location of expression in the plant. For SSS III the N-terminal extension is 780 amino acids long, for gbSS II 275. The C-terminal part of these two enzymes, SSS I, and GBSS I is very similar. Three regions in particular,
15 the putative ADP-glucose binding site. The function of the other boxes is unknown. GBSS I and gbSS II are expressed in stolon and tuber, SSS I only in leaf, and SSS III in both tuber and leaf. The amino acid sequence of potato BE I and BE II is highly conserved, except at the extremities of the protein. BE II has a flexible N-terminal extension of approximately 120 amino acids which is absent in BE I. BE I has a C-
20 terminal extension of 110 amino acids which BE II lacks. Contrary to maize, only two isoforms of BE have been found in potato so far. It should be noted that BE I is much more abundant than BE II in potato tubers. In maize three isoforms have been reported (BE I, IIa, and IIb). The isoforms BE IIa and BE IIb of maize are very similar with respect to their amino acid sequence (except for the c. 50 N-terminal amino acids).
25 However, these isoforms seem to be expressed in different tissues.
- Although the primary structure of many synthases and BEs has been documented, this is not the case for the biochemical properties of these enzymes. Based on observations in certain mutant backgrounds, there is some evidence suggesting that each synthase plays a particular role in elongating sidechains of a specific length.
30 Further, GBSS I is the only synthase involved in the synthesis of amylose. Next to (slightly) different catalytic properties, it is expected that the ability of synthases to bind to starch granules is determinative for their role in the biosynthesis process. GBSS I is found predominantly in the starch granule, whereas SSS III is found predominantly

11-06-1999

AGENINGEN

EP99201862.2

DESC

28

in the soluble phase. The gbSS II and SSS I seem to hold intermediate positions. The isoforms of BE seem to display a somewhat different mode of action. BE II prefers to transfer shorter glucan chains to an acceptor substrate than BE I. In addition, other properties of the BEs might determine their specific roles in the biosynthesis process.

- 5 For instance, BE II seems to associate more strongly with starch granules than BE I. In fact, one might say that BE II is partially granule-bound (maybe gbBE would be a more appropriate name). It is unknown whether its starch-binding ability is mediated through the N-terminal extension (which is typical for BE II).

- Over the years, the level of several of the synthase or BE isoforms has been
10 down-regulated by mutation or antisense technology. This large amount of data suggests that decreasing the level of granule-bound enzymes has more severe consequences for the starch granule architecture than that of their soluble isoforms. The presence of GBSS I determines whether or not amylose is deposited in starch granules. Knocking out SSS III expression in potato leads to a different granule morphology (T-
15 shaped cracks and clusters of small granules). However, starch content of the tubers, the granule size, and the ratio of amylopectin and amylose remain more or less unaltered. Inhibition of gbSS II expression in potato does not result in large changes in starch content, morphology or composition. However, mutation of the pea gbSS II gene can result in an altered granule morphology and amylopectin structure.

- 20 The impact of downregulating a particular synthase may be related to how much this isoform contributes to the total amount of synthase activity in the crop. However, this may be difficult to assess. BE I is the most abundant BE isoform in potato tuber, and it resides predominantly in the soluble phase. Antisensing the gene encoding this enzyme effectively removes the BE activity from the tuber juice, but surprisingly, the
25 starch structure is not affected by this. However, decreasing the level of BE II protein (which is at least partially granule-bound) in potato tuber by antisense technology dramatically reduces the number of α -(1 \rightarrow 6) branchpoints in the starch (and probably also the starch content of the tuber), and consequently the functional properties of the starch. The starch produced in this way shares some characteristics with the amylose-
30 extender mutant in maize, in which the gene encoding BE IIb is inactivated.

According to the invention, soluble enzymes could be made granule-bound; and techniques can be provided make other (non-)biosynthetic enzymes granule-bound.

11-06-1999

WAGENING

EP99201862.2

DESC

Protein-carbohydrate interactions are not understood with respect to starch biosynthetic enzymes (or granule-boundness), but the ability to bind to crystalline particles is a common feature of many cellulose- or starch-degrading enzymes. In general, these enzymes are composed of two or more domains which are connected by linker peptides. One of these domains contains the catalytic function of the protein, whereas other domains are involved in anchoring the proteins to the water-insoluble polysaccharide matrix. The so-called binding-domains (BDs) are usually relatively small (30 to approximately 160 amino acids) compared to the catalytic domains (>200 amino acids). As a result of this, the 3D structures of a large number of BDs have been solved. These studies have provided a rather detailed picture of how a protein interacts with carbohydrates. Further, a number of candidate amino acid residues which are presumably involved in binding the polysaccharide could be indicated. In some cases the importance of these residues for binding was verified by site-directed mutagenesis experiments. The interaction of cellulose-binding domains (CBDs) with cellulose and of (raw) starch-binding domains (SBDs) with starch granules will be further discussed hereinbelow.

Unlike CBDs, the amino acid sequences of SBDs seem very well conserved among different enzymes (α -amylase, β -amylase, glucoamylase, cyclodextrin glycosyltransferase [CGTase], etc.), as well as among different species (*Aspergillus niger*, *Bacillus circulans*, *Streptomyces limosus*, *Clostridium thermosulfurogenes*, *Pseudomonas stutzeri*, *Klebsiella pneumoniae*, etc.). In analogy to CBD type I, II, III, and V, SBDs are rather rigid structures which are predominantly composed of β -strands. Based on their interaction with maltose or cyclodextrin, usually two separate sugar-binding sites (1 and 2) can be distinguished in SBDs, which contain two or three exposed aromatic amino acids. The structure of site 1 is better conserved among the different SBDs than that of site 2, and has presumably a higher affinity for ligands than site 2. It contains two easily accessible Trp residues, which more or less keep their orientation upon ligand binding. Site 2 is much longer than site 1 and contains two or three Tyr residues which are located on a rather flexible loop of the SBD. Upon binding, site 2 undergoes significant structural changes which may allow the SBD to interact with starch in various orientations. In CGTases the SBD is part of a complex structure which comprises 5 separate domains. In these enzymes, site 1 is on the outside of the protein, whereas site 2 is more buried in the protein structure forming

11-06-1999

AGENINGEN

EP99201862.2

DESC

30

part of a channel leading to the catalytic site of the enzyme. Site-directed mutagenesis of aromatic amino acid residues belonging to both site 1 and 2 suggest that site 1 is the actual (raw) starch binding site whereas site 2 is involved in guiding glucan chains to the catalytic site of the CGTase. It is not known whether site 2 also contributes to

5 binding of starch granules.

Next to the SBDs, there are also proteins which only bind soluble α -glucans. An example of this is the maltodextrin-binding protein (MBP) from *Escherichia coli*. The amino acid residues that are involved in ligand binding are buried in the MBP structure. From the above, it can be seen that at least some the following factors may influence

10 protein-carbohydrate interactions. (i) Aromatic amino acids (often two or more) play a major role in these interactions. (ii) For binding crystalline structures these residues are positioned on an exposed face of the protein. (iii) For binding water-soluble molecules or amorphous structures these residues, or residues with a similar function are lining a groove, buried in the protein.

15 A number of enzymes involved in starch biosynthesis have been shown to associate with starch granules; one preferred subclass thereof according to the invention are the synthases. Reasons for this are the availability of several synthase genes and a mutant GBSS I (amf) potato plant (producing an amylose-free starch).

20 Example I: Determining factors involved in the granule-boundness of GBSS I.

WT, mutant and fusion proteins can be expressed in *E. coli* using a suitable expression system. Subsequently, the various enzymes can be purified, and their biochemical characteristics (activity, starch-binding) can be determined. Potato plants (both WT and amf background) can then be transformed with selected genes, based on

25 the properties of their corresponding proteins. The forthcoming transgenic starch granules can then be analyzed for alterations in their size, morphology, and composition (including the fine structure of the polysaccharides if necessary). Thus, polymer production can be studied *in planta* with well-characterized proteins providing the essentials to understand why this polymer was made.

30 The amino acid sequence of GBSSI is determined in a manner known per se, for instance from the nucleotide sequence of an isolated GBSSI-encoding cDNA. Based on amino acid sequence alignments of various synthases a number of Trp or Tyr residues in the GBSSI sequence can be selected as candidates for the interaction with starch

granules. These (and combinations of these) can be replaced by site-directed mutagenesis. The mutant proteins are then be purified and analyzed for a reduced starch-binding capacity.

Potato (amf background) is then transformed with mutant GBSS I with reduced affinity to determine whether amylose biosynthesis can be restored to a similar extent as with WT GBSS I.

Based on the above experiments, aromatic amino acids can be introduced at appropriate positions in *glgA* from *Bacillus subtilis*, to study whether this enzyme can be made granule-bound. Potato plants (amf background) can then be transformed with mutant *glgA* (s) to study whether amylose biosynthesis can be restored. In this way the starch-binding site of GBSS I can be mapped. Alternatively, the Tyr residues involved in binding could be replaced by Trp residues in order to tailor a GBSS I with improved starch-binding characteristics (as Trp residues are known to bind with higher affinity than Tyr residues).

Example II: Tailoring granule-boundness.

Granule-boundness may be tailored by fusing a relatively small SBD (100 amino acids), such as the SBD of the CGTase from *Bacillus circulans*, to a ("soluble") protein of interest.

This protein used can for instance be a soluble starch synthase, in order to provide amf potato plants that can be used as a model system for studying the potential of "artificial" granule-boundness *in planta* (i.e. to determine whether amylose biosynthesis in an amf background can be restored by expression of a fusion of the invention). Different fusions can be made. Next to synthases equipped with one SBD, also synthases with a two SBDs can be expressed, optionally with variations in in the linker peptides connecting synthase and SBD(s). The fusion proteins can also be expressed in *E. coli*, and subsequently purified to investigate their biochemical properties (starch-binding, activity). Potato plants (amf background) can be transformed with selected fusion proteins to study whether amylose biosynthesis can be restored to a similar extent as with WT GBSS I.

In addition, WT potato plants can be transformed with with a gene encoding a single SBD or a double SBD to determine the competition between GBSS I and

SBD(s) *in vivo*. This may also as an alternative way to make an amylose-free potato starch.

In these experiments, transformation of WT potato plants with (double) SBDs will also show whether GBSS I and SBD bind to similar parts (or structures) in the starch granule (i.e whether GBSS I and SBD differ in their binding specificity). A different binding specificity would also provide the opportunity to target proteins to different parts of the starch granule, and also to use an inactive GBSS I (for instance obtained via genetical modification) as an alternative for a SBD in fusions. In addition, by making deletion mutants, different parts of GBSS I can be tested and used for their affinity for starch. (In this manner, alternative starch-binding domains could be obtained, with the same binding properties as GBSS I, but (much) of smaller size and and without catalytic activity.)

Example III: Enzyme fusions and possible applications.

Three possible enzymes that can be equipped with an SBD by expression as a fusion of the invention are the *Escherichia coli* glycogen branching enzyme glgB, the potato kinase R1, and a glucose oxidase/dehydrogenase. The former two have already been expressed in potato without a SBD. By expression of these proteins as a fusion of the invention, their impact on starch structure and functionality can be increased.

Example III-1: Preparation of extra heavily branched amylopectin.

In a previous investigation by Applicant, the amount of branching in starch polymers was increased by introducing a heterologous branching enzyme in both WT and amf potato plants. The outcome of these experiments was as follows. (i) In an amf background an additional 25% of branching of the amylopectin was obtained. (ii) In a WT background evidence was obtained that next to the extra branching of amylopectin, amylose disappears as such. (Assuming that amylose is synthesized downstream of amylopectin, this indicates that glgB is (partially) granule-bound.)

Additional branching comparwed to these reference tests might be achieved when glgB is specifically targeted to (and thus concentrated at) the granule surface by equipping the enzyme with a SBD, i.e. by expressing it *in planta* as a fusion of the invention. In this way, with a fusion of the invention, possibly a larger increase in

11-06-1999

EP99201862.2

DESC

33

branching can be obtained (i.e. compared to transforming with glgB per se), to provide improved freeze-thaw stability of starch solutions.

Example III-2: Phosphorylated starch.

5 Potato starch differs from many other starches in that it is more heavily phosphorylated. There is roughly one phosphate group to 700 Glc residues. Phosphorylation can occur at the C-6 (65%) or the C-3 position (35%) of Glc. The relatively large phosphate content gives potato tuber starch a number of unique properties such as the "peak viscosity" upon gelatinizing the granule, an anionic
10 character, and possible anchors for derivatization.

Recently, an enzyme has been cloned which is presumably responsible for phosphorylation of starch. This kinase, further referred to as R1, cleaves ATP, and transfers a phosphate group to an unknown donor molecule. Phosphorylation of starch could be further increased by expressing a fusion of an SBD and R1 *in planta*, for
15 instance in potato or cassava (heterologous expression).

Example III-3: Oxidized starch.

Starch contains a large collection of hydroxyl groups ($-\text{[CH}_2\text{]-OH}$), but these are not very reactive. Next to these, each starch polymer contains an aldehyde group
20 ($-\text{[CH]=O}$) at the reducing terminus, which is much more reactive. Because the starch polymers are such large molecules, the number of reactive groups is too small to be a meaningful target for derivatization. One way to increase the number of aldehyde or carboxyl ($-\text{[COH]=O}$) groups is by oxidation. As an alternative to chemical oxidation procedures, oxidases or dehydrogenases can be used, i.e. by expression as a fusion
25 according to the invention. This could already lead to (increased) oxidation *in planta*, but also, and preferably, an oxidative enzyme fused to an SBD is incorporated in a starch granule during granule biogenesis, and subsequently these granules are incubated *in vitro* under conditions (temperature, pH, co-factors) suitable to provide oxidation of the starch.

30

Example IV: Constructs of the invention.

Schematic representations of some non-limiting examples of constructs of the invention are shown in Figures 2A – E (in which TP represents a leader sequence, i.e.

11-06-1999

EP99201862.2

DESC

34

the transit peptide of potato GBSSI). Instead of the GUS-gene shown in Figure 2, also another reporter gene such as (a sequence encoding a) luciferase can be used, or a sequence encoding the desired protein or polypeptide.

11-06-1999

EP99201862.2

EPO - DG 1

CLMS

11. 06. 1999

C L A I M S

(90)

1. Method for expressing a desired protein or polypeptide in a plant, in which the
5 protein or polypeptide is expressed as a fusion with at least one starch binding domain.

2. Method according to claim 1, comprising the steps of:

- 10 a) providing a genetic construct comprising at least one nucleotide sequence encoding the
desired protein or polypeptide combined with at least one nucleotide sequence encoding a
starch binding domain, so that the construct encodes a fusion of the desired
protein/polypeptide and the at least one starch binding domain;
b) transforming a plant with said genetic construct;
c) expressing said genetic construct in the plant.

15 3. Method according to claim 1 or 2, in which the plant is a plant that contains or
produces starch or starch granules in at least one of its parts, including its seeds, leaves, roots
(including tuberos roots), tubers, stems, stalks, fruits, grains or flowers (in particular the
honey-producing parts thereof).

20 4. Method according any of the preceding claims, in which the plant is chosen from
potato, sweet potato, cassava, pea, taro, sago, yam, banana and/or cereals such as rice, maize,
wheat and barley.

25 5. Method according to any of the preceding claims, in which the protein or polypeptide
is heterologous with respect to the plant in which the fusion is expressed.

6. Method according any of the preceding claims, in which the protein or polypeptide is
an enzyme.

30 7. Method according to claim 6, in which the enzyme is an enzyme that can interact
with starch or starch granules, in particular an enzyme that can convert, modify, alter, degrade
or otherwise influence the starch, the starch granule or the structure or interactions thereof.

11-06-1999

EP99201862.2

CLMS

8. Method according to any of claims 1-5, in which the desired protein or polypeptide is a receptor such as an estrogen receptor or a plant hormone receptor, or a structural protein such as a protein "zipper".

9. Fusion of at least one desired protein or polypeptide and at least one starch binding domain, as expressed in and/or present in a plant or in any part of a plant, including its seeds, leaves, roots (including tuburous roots), tubers, stems, stalks, fruits, grains or flowers (in particular the honey-producing parts thereof).

10. Genetic construct suitable for transforming a plant, comprising at least one nucleotide sequence encoding a desired protein or polypeptide combined with at least one nucleotide sequence encoding a starch binding domain, so that the construct encodes a fusion of the desired protein/polypeptide and the at least one starch binding domain.

11. Method for providing a plant that expresses a fusion according to claim 9, comprising at least one step of:

- a) transforming plant with a genetic construct according to claim 10, such that said genetic construct is expressed in the plant or at least in part thereof; and optionally further comprising at least one step of:
- b) providing descendants and/or further generations of the thus transformed plant, for instance via sexual or asexual multiplication, including crossing and/or other breeding techniques.

12. Method according to claim 11, in which the plant is a plant that contains or produces starch or starch granules in at least one of its parts, including its seeds, leaves, roots (including tuburous roots), tubers, stems, stalks, fruits, grains or flowers (in particular the honey-producing parts thereof).

13. Plant, transformed with a genetic construct according to claim 10, or a descendant of such a plant.

14. Plant that expresses a fusion according to claim 9, obtainable by the method of claim 10.

11-06-1999

EP99201862.2

15. Seeds, tubers, seedlings or other cultivating material of a plant according to claim 13 or 14.

16. Method for providing a protein or polypeptide with affinity for starch and/or starch granules, and/or for increasing the affinity of a protein or polypeptide for starch and/or starch granules, comprising expressing the protein or polypeptide in a plant as a fusion with at least one starch binding domain.

17. Method according to claim 16, comprising the steps of:

- a) combining a nucleotide sequence encoding the protein or polypeptide with at least one nucleotide sequence encoding a starch binding domain, so as to provide a genetic construct encoding a fusion of the protein or polypeptide and the at least one starch binding domain;
- b) transforming a plant with said genetic construct;
- c) expressing said genetic construct in the plant.

18. Method for producing a complex of at least one protein or polypeptide and a starch granule, comprising at least one step of:

- a) expressing the protein or polypeptide as a fusion with at least one starch binding domain, in a plant that contains or forms starch granules; and optionally comprising a further step of:
- b) isolating the protein or polypeptide from the plant or any part thereof as a complex of the fusion and the starch granule.

19. Complex, comprising a fusion of a protein or polypeptide fused to at least one starch binding domain, associated with a starch granule.

20. Method for providing a plant that can produce a complex according to claim 19, comprising at least the step of:

- a) transforming a starch granule producing plant with a genetic construct according to claim 19, such that said genetic construct is expressed in the plant or at least part thereof; and optionally further comprising at least a further step of:

11-06-1999

EP99201862.2

CLMS

- b) providing descendants and/or further generations of the thus transformed plant, for instance via sexual or asexual multiplication, including crossing and/or other breeding techniques.

5 21. Plant that produces a complex according to claim 19, and that is obtainable via the method of claim 20, or any descendant thereof.

22. Seeds, tubers, seedlings or other cultivating material of a plant according to claim 21.

10 23. Method for providing a plant that contains or produces a modified starch and/or modified starch granules, comprising at least one step of:

- a) transforming a starch producing plant, in particular a starch granules producing plant, with a genetic construct comprising at least one nucleotide sequence encoding an enzyme that can interact with starch and/or starch granules and at least one nucleotide sequence
15 encoding a starch binding domain, such that said genetic construct is expressed in the plant or at least part thereof;

and optionally further comprising at least a further step of:

- b) providing descendants and/or further generations of the thus transformed plant, for instance via sexual or asexual multiplication, including crossing and/or other breeding
20 techniques.

24. Plant that produces modified starch or starch granules, and that is obtainable via the method of claim 23, or any descendant thereof.

25 25. Seeds, tubers, seedlings or other cultivating material of a plant according to claim 24.

26. Method for producing a modified starch and/or modified starch granules, comprising at least one step of:

- a) cultivating a transformed plant that produces a modified starch and/or modified starch
30 granules according to claim 24, or a descendant thereof;
and optionally further comprises at least one step of:
b) isolating the modified starch or starch granules from the transformed plant or any part thereof, including its seeds, leaves, roots (including tuberos roots), tubers, stems, stalks, fruits, grains or flowers (in particular the honey-producing parts thereof).

11-06-1999

EP99201862.2

CLMS

27. Method for producing a modified starch and/or modified starch granules, comprising at least one step of:

- 5 a) cultivating a transformed plant that expresses a fusion of at least one starch binding domain and at least one enzyme that can interact with starch (granules), in which the starch (granule) and the fusion preferably form a complex;
- b) harvesting the plant and/or any part of said plant that contains starch (granules) and the fusion, such as the seeds, leaves, roots (including tuberos roots), tubers, stems, stalks, fruits, grains or flowers (in particular the honey-producing parts thereof);
- 10 c) subjecting the plant or plant material, or any fraction or preparation obtained therefrom that contains the starch (granule) and the fusion, to conditions such that the enzyme can interact with the starch (granule) to provide modified starch or starch granules; and optionally comprises the further step of:
- d) isolating the modified starch or starch granules thus obtained.

15

28. Modified starch or starch granules, obtained via the method of claim 26, obtained from a plant according to claim 24 or any part thereof, or obtained via the method of claim 27.

29. Method for providing a plant that contains or produces a modified starch and/or modified starch granules, comprising at least one step of:

- 20 a) transforming a starch producing plant, in particular a starch granules producing plant, with a genetic construct comprising at least one nucleotide sequence encoding a starch binding domain, such that said genetic construct is expressed in the plant or at least part thereof; and optionally further comprising at least one step of:
- 25 b) providing descendants and/or further generations of the thus transformed plant, for instance via sexual or asexual multiplication, including crossing and/or other breeding techniques.

30. Plant that produces modified starch or starch granules, obtainable via the method of claim 29, or any descendant thereof.

30

31. Seeds, tubers, seedlings or other cultivating material of a plant according to claim 30.

11-06-1999

EP99201862.2

CLMS

32. Method for producing a modified starch and/or modified starch granules, comprising at least one step of:~

a) cultivating a transformed plant that produces a modified starch and/or modified starch granules according to claim 30, or a descendant thereof;

5 and optionally further comprises at least one step of:

b) isolating the modified starch or starch granules from the transformed plant or from any part thereof, including its seeds, leaves, roots (including tuberos roots), tubers, stems, stalks, fruits, grains or flowers (in particular the honey-producing parts thereof).

10 33. Modified starch or starch granules, obtained via the method of claim 32, or obtained from a plant according to claim 30 or any part thereof.

34. Modified starch according to claim 33, being an essentially amylose-free starch or starch granule.

15

35. Genetic construct suitable for transforming a plant, comprising at least one nucleotide sequence encoding a starch binding domain.

20 36. Bacterium, virus or other organism suitable for transforming a plant, containing a genetic construct according to claim 10 or 35, and preferably capable of transferring said construct into a plant.

11-06-1999

WAGENINGEN

EP99201862.2

PO - DG 1

ABST

ABSTRACT

11. 06. 1999

(90)

The invention relates to a method for expressing a desired protein or polypeptide in a plant, in which the protein or polypeptide is expressed as a fusion with at least one starch binding domain.

The plant is preferably a plant that contains or produces starch or starch granules in at least one of its parts, such as potato, sweet potato, cassava, pea, taro, sago, yam, banana and/or cereals such as rice, maize, wheat and barley.

The protein or polypeptide can be an enzyme, in particular an enzyme that can convert, modify, alter, degrade or otherwise influence starch (granules); or can be a receptor or a structural protein.


The invention further relates to the fusions thus obtained, to genetic constructs that encode the above fusions and to plants transformed with said constructs.

The method of the invention can in particular be used to provide modified starches and/or to provide complexes of starch (granules) and the above fusions.

In another embodiment, one or more starch binding domains are expressed in a plant, to provide a plant producing modified starches.



Fig. 1: Alignment of starch-binding domains



■ maltose-binding site 1 · ■ maltose-binding site 2 ■ disulfide bridge

DRAW


\\SBD (4x) alignment

11-06-1999


Fig. 2:

Constructs for targeting proteins to starch granules


2A: SBD




2B: SBD2



2C: SBD-GUS



2D: GUS-SBD



2E: GBSS I

